maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to completing and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding ar OMB control number.	ion of information. Send comments arters Services, Directorate for Info	s regarding this burden estimate or ormation Operations and Reports	or any other aspect of the 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington	
. REPORT DATE 2007		2. REPORT TYPE		3. DATES COVERED 00-00-2007 to 00-00-2007		
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER		
Helical Magnetic Fields in the Jet of 3C273				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAIL Approved for publ	LABILITY STATEMENT ic release; distributi	on unlimited				
13. SUPPLEMENTARY NO	OTES					
	e confirm the present on measure gradient stic jet.					
16. SECURITY CLASSIFICATION OF: 17. LIMITATION OF:				18. NUMBER	19a, NAME OF	
			ABSTRACT	OF PAGES	RESPONSIBLE PERSON	
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	2		

Report Documentation Page

Form Approved OMB No. 0704-0188

Helical Magnetic Fields in the Jet of 3C 273

R. T. Zavala

United States Naval Observatory

G. B. Taylor

Department of Physics and Astronomy, University of New Mexico

G. Giovannini

INAF-Instituo di Radioastronomia and Univ. Bologna

Abstract. Using the VLBA we confirm the presence of a Faraday rotation measure gradient transverse to the jet axis of 3C 273. A rotation measure gradient is expected to be the signature of a helical magnetic field wrapping around the relativistic jet.


1. The Case for Helical Magnetic Fields in 3C 273

The role of magnetic fields in relativistic jets in AGN is a subject of considerable interest. Simulations have produced results that suggest that a helical magnetic field may wind around the jet (Meieret al. 2001; Koide et al. 2002). Before these simulations took place Blandford (1993) suggested that observers look for evidence of a rotation measure (RM) gradient transverse to the jet axis. Such a gradient is expected if a helical or toroidal field is present in the jet and if such a field embedded in a plasma can act as a Faraday rotating screen. Asada et al. (2002) reported the detection of such an RM gradient transverse to the jet axis in 3C 273. These 5 and 8 GHz VLBA observations just began to resolve the jet. We examined higher frequency data and confirmed the presence of the RM gradient over two epochs separated by six months (Zavala & Taylor 2005). We showed that the transverse RM gradient persists over two epochs separated by 6 months. In our poster we reproduced Figures 1 and 3 from Zavala & Taylor (2005). The sense of RM gradient detected in Asada et al. (2002) and Zavala & Taylor (2005) was confirmed by Attridge et al. (2005) at less than 3 pc from the core. This gradient appears to change in magnitude as one proceeds along the jet (Wardle et al. 2006).

This result led us to search for other sources that may exhibit an RM gradient transverse to the jet axis. Other investigators have reported such gradients. However, the amplitude of the gradient is less than that observed in 3C 273 (e.g., Gabuzda et al. 2004). We observed four sources that have broad, polarized jets with the VLBA. Here we present preliminary results for the blazar Mrk 501.

2. Mrk 501

Our VLBA observation of Mrk 501 used frequencies from 8–22 GHz. The RM map is in Fig. 1, and the black line indicates the cut along which we plot the RM in the right panel. No significant gradient in the RM transverse to the jet






Figure 1. Left panel shows the RM map of Mrk 501 overlaid on total intensity contours. The black line represents the slice across the RM distribution. The slice is shown in the right panel as Kilo $\operatorname{rad/m^2}$ and mas along the slice.

is seen in Mrk 501. As Mrk 501 is a TeV blazar the jet we observe may be significantly different from that in 3C 273. A blazar jet more aligned to our line-of-sight may produce very different Faraday rotation effects from that seen in a more inclined quasar jet. A blazar jet could clear out some of the thermal gas of a Faraday screen. At the scale of Fig. 1 the jet in Mrk 501 changes its orientation angle significantly, which may complicate the picture we see in the RM maps (Giroletti et al. 2004). However, Croke et al. (2005) report an RM gradient suggestive of a helical B field from earlier VLBA epochs.

Acknowledgments. R.T.Z. received an International Travel Grant from the American Astronomical Society.

References

Attridge, J. M., Wardle, J. F. C., & Homan, D. C. 2005, ApJ, 633, L85 Asada, K., et al. 2002, PASJ, 54, L39

Blandford, R. D. 1993, in Astrophysical Jets, ed. D. Burgarella, M. Livio & C. P. O'Dea (Cambridge: Cambridge University Press), 15

Croke, S., Charlot, P., Gabuzda, D., & Sol, H. 2005, Baltic Astronomy, 14, 367

Gabuzda, D. C., Murray, É., & Cronin, P. 2004, MNRAS, 351, L89

Giroletti, M., et al. 2004, ApJ, 600, 127

Koide, S., et al. 2002, Science, 295, 1688

Meier, D. L., Koide, S., & Uchida, Y. 2001, Science, 291, 84

Wardle, J. et al. 2006, in online proceedings of the conference Challenges of Relativisitc Jets, ed. M. Ostrowski & L. Stawarz (http://www.oa.uj.edu.pl/2006jets)

Zavala, R. T., & Taylor, G. B. 2005, ApJ, 626, L73